

buruli ulcer mycobacterium ulcerans disease Updated Version

Buruli ulcer mycobacterium ulcerans disease is a debilitating skin infection caused by the bacterium *Mycobacterium ulcerans*. This disease predominantly affects populations in tropical and subtropical regions, especially in West Africa, Central Africa, and some parts of Australia, Asia, and the Americas. Characterized by painless skin ulcers, nodules, and plaques, Buruli ulcer can lead to severe tissue destruction if not diagnosed and treated early. Understanding the causes, symptoms, transmission, diagnosis, and treatment options for this disease is crucial for public health efforts and affected communities.

Understanding Buruli Ulcer Mycobacterium Ulcerans Disease

Buruli ulcer is classified as a neglected tropical disease, often overshadowed by more prominent infections but nonetheless posing a significant health burden in endemic regions. The causative agent, *Mycobacterium ulcerans*, is a slow-growing bacterium related to the bacteria responsible for tuberculosis and leprosy. Its unique pathogenicity is primarily due to the production of a toxin called mycolactone, which causes tissue necrosis and immune suppression.

Causes and Transmission of Buruli Ulcer

The Role of Mycobacterium ulcerans

Mycobacterium ulcerans is a slow-growing mycobacterium that thrives in aquatic environments. It is believed to be transmitted through contact with contaminated water or environmental reservoirs. The bacteria can enter the skin through cuts, abrasions, or insect bites, leading to localized infection.

How Is Buruli Ulcer Transmitted?

The exact mode of transmission remains an area of ongoing research, but several pathways have been identified:

- Environmental Exposure: Contact with contaminated water bodies such as lakes, swamps, and slow-moving rivers is a primary risk factor.
- Insect Vectors: Some studies suggest that aquatic insects or mosquitoes may play a role in transmitting the bacteria.
- Soil and Vegetation: Contact with soil or vegetation contaminated with *Mycobacterium ulcerans*

may also contribute to infection.

Key Points About Transmission:

- The disease is not directly contagious from person to person.
- Environmental factors, including warm, humid climates, facilitate bacterial survival.
- People involved in farming, fishing, or living near water bodies are at higher risk.

Symptoms and Clinical Features of Buruli Ulcer

Early Signs

Initially, the infection may present as:

- A painless nodule or swelling under the skin
- A flat, bluish or purple plaque
- A thickened skin area or raised lesion

These early signs often go unnoticed due to the painless nature of initial lesions.

Progression of the Disease

Without treatment, the lesion enlarges over weeks or months, leading to:

- Ulceration with undermined edges
- Extensive destruction of skin, soft tissue, and sometimes bones
- Possible secondary bacterial infections

Characteristic Features:

- Usually painless, which can delay seeking medical help
- Often occurs on the limbs, especially the arms and legs
- Can cause significant scarring and disfigurement

Diagnosis of Buruli Ulcer

Accurate and early diagnosis is essential for effective management. Diagnostic methods include:

- Clinical Evaluation: Recognizing characteristic skin lesions and swelling
- Laboratory Tests:
 - Polymerase Chain Reaction (PCR): Highly sensitive for detecting Mycobacterium ulcerans DNA
 - Microscopy: Acid-fast bacilli staining
 - Culture: Growing the bacteria, though it takes several weeks
 - Histopathology: Tissue biopsy showing necrosis and acid-fast bacilli
 - Imaging: Ultrasound or MRI may assess tissue extent

Challenges in Diagnosis:

- Similar appearance to other skin conditions
- Limited laboratory facilities in endemic regions
- Need for trained healthcare professionals

Prevention and Control Strategies

Preventing Buruli ulcer involves reducing exposure to environmental sources and early detection. Key preventive measures include:

- Avoiding contact with contaminated water bodies, especially during activities like bathing or fishing
- Using protective clothing and footwear
- Promoting community awareness about early signs and symptoms
- Enhancing sanitation and water management

Community-Based Approaches:

- Educating at-risk populations
- Implementing active case detection programs
- Providing access to healthcare facilities for early diagnosis

Treatment Options for Buruli Ulcer

Effective management of Buruli ulcer revolves around antibiotics, surgical intervention, and supportive care.

Antibiotic Therapy

The World Health Organization recommends a combination of antibiotics for at least 8 weeks:

- Primary Regimen:
- Rifampicin plus streptomycin (for small to moderate lesions)
- Rifampicin plus clarithromycin (alternative for some cases)
- Outcome: Antibiotics help halt bacterial growth, prevent further tissue destruction, and promote healing.

Surgical Intervention

In some cases, surgery is necessary to:

- Excise necrotic tissue
- Debride ulcers
- Reconstruct affected areas after infection control

Supportive and Wound Care

- Proper wound management to prevent secondary infections
- Pain management
- Nutritional support to aid tissue repair

Prognosis and Outcomes

With early diagnosis and appropriate treatment, most patients recover fully without significant disability. However, delays in treatment can lead to:

- Extensive tissue destruction
- Bone involvement
- Permanent scars and deformities
- Psychological and social impacts due to disfigurement

Research and Advances in Buruli Ulcer Management

Ongoing research aims to:

- Develop faster diagnostic tools
- Explore new antibiotics and treatment regimens
- Understand the ecology and transmission pathways better
- Create effective vaccines

Recent advances include the development of point-of-care diagnostic tests and shorter antibiotic regimens, which could improve patient compliance and outcomes.

Global Buruli Ulcer Disease Burden

Buruli ulcer remains a neglected disease with significant public health implications:

- Estimated thousands of new cases annually
- Endemic regions primarily in Africa, with emerging cases worldwide
- Limited awareness and healthcare infrastructure in some endemic zones

Efforts by organizations such as WHO, CDC, and local health authorities focus on surveillance, education, and resource allocation to control the disease.

Conclusion

Buruli ulcer mycobacterium ulcerans disease is a complex skin infection that poses a significant health threat in tropical and subtropical regions. Its insidious onset, painless lesions, and potential for disfigurement underscore the importance of early detection and treatment. Preventive strategies centered around reducing environmental exposure, community education, and strengthening healthcare systems are vital. Advances in diagnostics, treatment protocols, and research hold promise for better management and eventual eradication of Buruli ulcer globally. Public awareness and international cooperation are essential to combat this neglected tropical disease effectively.

Key Takeaways:

- Buruli ulcer is caused by *Mycobacterium ulcerans*, transmitted mainly through environmental exposure.
- Early signs are painless skin nodules or plaques that can progress to ulcers.
- Diagnosis involves clinical evaluation and laboratory tests like PCR.
- Treatment primarily involves antibiotics, with possible surgical intervention.
- Prevention focuses on reducing contact with contaminated water and early case detection.
- Continued research is crucial for improved diagnostics, treatment, and understanding of transmission.

By understanding the disease comprehensively, affected populations and healthcare providers can work together to reduce the burden of Buruli ulcer and improve patient outcomes worldwide.

Buruli ulcer Mycobacterium ulcerans disease: An in-depth review of a neglected tropical infection

Introduction

Buruli ulcer, caused by Mycobacterium ulcerans, is a chronic necrotizing disease that predominantly afflicts populations in tropical and subtropical regions. Recognized as one of the world's neglected tropical diseases by the World Health Organization (WHO), it leads to significant morbidity, disability, and social stigma among affected communities. Despite its long history of identification, much about its transmission, ecology, and optimal treatment strategies remains elusive, underscoring the need for continued research and public health interventions.

Understanding Buruli Ulcer: An Overview

What is Buruli Ulcer?

Buruli ulcer is characterized by the development of painless skin lesions, often starting as a small nodule or papule, which can progress to large ulcers with undermined edges. The disease primarily affects the skin and underlying soft tissues, leading to extensive tissue destruction if left untreated. It predominantly occurs on the limbs, especially the arms and legs, but can also affect other parts of the body.

The causative agent, Mycobacterium ulcerans, is a slow-growing environmental mycobacterium that produces a potent toxin called mycolactone. This toxin is central to the pathogenesis of the disease, causing tissue necrosis and immunosuppression, which complicate diagnosis and treatment.

Epidemiology and Geographic Distribution

Global Distribution and Buruli Ulcer Hotspots

Buruli ulcer is endemic in more than 30 countries across Africa, Asia, the Americas, and the Western Pacific. The highest burden is observed in West and Central Africa, with Nigeria, Cote d'Ivoire, and Ghana among the most affected nations. Other notable regions include Australia, particularly in southeastern Queensland and Victoria, and parts of Southeast Asia such as Malaysia and Thailand.

Key epidemiological features include:

- Age distribution: It affects all age groups but is most common among children under 15.
- Gender distribution: Slightly more prevalent in males, possibly due to occupational or environmental exposures.
- Seasonality: Cases often surge during the rainy season, correlating with increased environmental activity of the bacteria.

Despite its widespread presence, the true incidence of Buruli ulcer is believed to be underestimated due to diagnostic challenges and underreporting.

Transmission Dynamics: Unraveling the Ecological Mysteries

Understanding How Mycobacterium ulcerans Spreads

One of the most intriguing aspects of Buruli ulcer is its elusive mode of transmission. Unlike many infectious diseases, direct human-to-human transmission appears to be rare or non-existent. Instead, the infection is thought to originate from environmental sources, making ecological and behavioral factors critical in disease dynamics.

Potential transmission pathways include:

- Environmental reservoirs: The bacteria are found in aquatic environments such as slow-moving or stagnant waters, swampy areas, and wetlands. The presence of *M. ulcerans* DNA has been detected in water, mud, and biofilms.
- Aquatic insects: Some studies suggest that certain aquatic insects, like water bugs (e.g., *Naucoris* species), can harbor the bacteria and potentially transmit it to humans through bites or skin breaches.
- Soil and vegetation: Contamination of soil and plant material in endemic areas may serve as environmental reservoirs.
- Human activities: Water contact during fishing, bathing, farming, or other recreational activities increases exposure risk.

Challenges in understanding transmission include:

- The bacteria's slow growth rate and environmental persistence.
- Difficulty in culturing *M. ulcerans* from environmental samples.
- The complex ecology involving multiple potential reservoirs and vectors.

Research efforts continue to clarify the precise mechanisms, which are pivotal for developing effective prevention strategies.

Pathogenesis: How Mycobacterium ulcerans Causes Disease

The Role of Mycolactone in Disease Development

The pathogenesis of Buruli ulcer hinges on the production of mycolactone, a lipid-like exotoxin that is both cytotoxic and immunosuppressive. This toxin is unique among pathogenic mycobacteria and is central to the disease's clinical features.

Mechanisms of pathogenicity include:

- Tissue necrosis: Mycolactone induces apoptosis and necrosis of skin and subcutaneous tissues, leading to painless ulcers.
- Immunomodulation: The toxin suppresses local immune responses, hindering the body's ability to contain the infection and facilitating bacterial proliferation.
- Delayed immune response: Patients often present with painless lesions because mycolactone diminishes pain and inflammation, delaying diagnosis.

Bacterial invasion begins with inoculation into the skin through minor trauma, after which *M. ulcerans* proliferates locally, secretes mycolactone, and causes progressive tissue destruction.

Clinical Presentation and Disease Progression

Signs, Symptoms, and Stages of Buruli Ulcer

The clinical course of Buruli ulcer can be divided into several stages:

- Initial Lesion: Usually a painless papule, nodule, or plaque. It is often unnoticed or mistaken for a benign skin condition.
- Ulceration: The lesion enlarges, with undermined edges, revealing a necrotic, ulcerative area. The ulcer is characteristically painless, which can delay seeking treatment.
- Advanced Disease: Without intervention, lesions may coalesce, cause extensive tissue destruction, and involve underlying structures such as muscles and bones, leading to deformity and disability.

Other clinical features include:

- Multiple lesions may occur in some cases.
- Edema and induration around the ulcer.

- Rarely, systemic symptoms like fever are observed, mainly in disseminated or severe cases.

Diagnosis in early stages is often missed due to the benign appearance and lack of pain, emphasizing the importance of awareness and early detection.

Diagnostic Strategies: Confirming the Disease

Laboratory and Clinical Diagnostic Tools

Accurate and timely diagnosis is crucial to prevent disfigurement. Diagnostic methods include:

- Microscopy: Acid-fast bacilli (AFB) detection using Ziehl-Neelsen staining can confirm *M. ulcerans* presence but has limited sensitivity.
- Culture: Growing *M. ulcerans* from clinical specimens is definitive but slow, requiring several weeks and specialized media.
- Polymerase Chain Reaction (PCR): The most sensitive and specific method, detecting *M. ulcerans* DNA, particularly targeting IS2404 insertion sequences.
- Histopathology: Biopsy of the lesion shows necrosis with acid-fast bacilli and characteristic infiltration patterns.
- Imaging: Ultrasound or MRI can assess the extent of tissue involvement.

Field-friendly diagnostics such as loop-mediated isothermal amplification (LAMP) assays are under development to facilitate point-of-care testing, especially in resource-limited settings.

Treatment Modalities: Combining Medical and Surgical Approaches

Current Treatment Strategies

Effective management of Buruli ulcer involves a combination of antibiotic therapy and, when necessary, surgical intervention.

Antibiotic Regimens:

- The WHO recommends an 8-week course of oral rifampicin combined with either streptomycin (for hospitalized cases) or clarithromycin.
- Recent studies have supported the efficacy of all-oral regimens, improving compliance.
- Antibiotics aim to sterilize the infection, prevent progression, and reduce transmission.

Surgical Management:

- Indicated for large ulcers, abscesses, or extensive tissue destruction.
- Procedures include debridement, excision, and skin grafting.
- Surgery is often combined with antibiotics for optimal outcomes.

Emerging therapies and adjunctive measures:

- Use of wound dressings and physiotherapy.
- Management of complications such as secondary infections.
- Early detection and treatment significantly reduce disability.

Challenges in Treatment:

- Delayed diagnosis leads to larger, more complex ulcers requiring extensive surgery.
- Antibiotic resistance, though currently rare, remains a concern.
- Limited access to healthcare facilities in endemic regions hampers timely intervention.

Prevention and Control: Strategies Towards Elimination

Public Health Measures and Future Directions

Prevention of Buruli ulcer focuses on reducing exposure and early detection.

Key strategies include:

- Community education: Raising awareness about the disease's signs and the importance of early treatment.
- Environmental management: Modifying water bodies and habitats to reduce bacterial reservoirs and vectors.
- Protective behaviors: Using protective clothing during water contact activities.
- Surveillance and reporting: Strengthening health systems for early case detection and response.

Research priorities encompass:

- Understanding environmental ecology and transmission pathways.
- Developing rapid, affordable point-of-care diagnostics.
- Exploring vaccine candidates, although none are currently available.
- Investigating the role of environmental interventions in disease reduction.

The Challenges and Future Outlook

Despite advances, significant gaps in understanding and controlling Buruli ulcer persist. The disease's obscure ecology, difficulties in early diagnosis, and limited access to treatment in endemic regions pose ongoing challenges. Climate change, environmental modifications, and human encroachment into natural habitats may influence disease epidemiology in unpredictable ways.

Future research must aim to:

- Clarify transmission dynamics.
- Develop effective vaccines and shorter, more tolerable treatment regimens.
- Implement integrated control strategies tailored to local contexts.
- Foster international collaboration and funding to address this neglected disease.

Conclusion:

Question What is Buruli ulcer and how is it caused? Buruli ulcer is a chronic skin disease caused by the bacterium *Mycobacterium ulcerans*. It leads to painless skin ulcers, primarily affecting people in tropical and subtropical regions. The bacteria produce a toxin called mycolactone, which destroys skin and soft tissue. **What are the common symptoms of Buruli ulcer?** Early symptoms include painless swelling or a nodule on the skin, often on limbs or the trunk. As the disease progresses, it causes large ulcers with undermined edges, which can lead to tissue destruction and disability if untreated. **How is Buruli ulcer diagnosed and treated?** Diagnosis typically involves clinical examination, histopathology, and laboratory tests such as PCR to detect *Mycobacterium ulcerans* DNA. Treatment generally includes an 8-week course of antibiotic therapy with drugs like rifampicin and streptomycin or clarithromycin, often combined with surgical removal of the affected tissue. **What are the risk factors for contracting Buruli ulcer?** Risk factors include residing in or visiting endemic areas, exposure to slow-moving or stagnant water bodies, and activities involving contact with contaminated aquatic environments. Poor hygiene and immunosuppression may also increase susceptibility. **Are there any preventive measures for Buruli ulcer?** Preventive measures focus on reducing contact with contaminated water sources, wearing protective clothing during water-related activities, and community education. Early detection and treatment are crucial to prevent severe disfigurement and complications.

Related keywords: Buruli ulcer, *Mycobacterium ulcerans*, skin infection, tropical disease, necrotizing ulcer, *Mycobacteria*, neglected tropical disease, antibiotic treatment, environmental reservoir, endemic regions

The Retreat Of Tuberculosis 1850 1950

The retreat of tuberculosis 1850 1950 marks a significant period in medical history characterized by remarkable progress in understanding, preventing, and treating one of humanity's oldest and most deadly infectious diseases. This era witnessed a transformation from widespread, often fatal, infections to a controlled disease with declining mortality rates, thanks to advancements in medical science, public health initiatives, and social reforms. Exploring this century-long journey offers insights into the scientific, social, and technological factors that contributed to this remarkable retreat.

Historical Context of Tuberculosis Before 1850

Pre-Industrial Era and Early Understanding

Before the mid-19th century, tuberculosis, often called "consumption," was a leading cause of death across Europe and North America. Its contagious nature was recognized, but the precise cause remained elusive, and treatments were largely ineffective. Factors such as poor living conditions, overcrowding, and malnutrition exacerbated its spread.

Social Impact and Stigma

Tuberculosis carried a heavy social stigma, often associated with poverty and moral failing. Patients were frequently isolated in sanatoria, which, despite their efforts, had limited success in controlling the disease's spread.

Scientific Breakthroughs and Advances (1850-1950)

Discovery of Mycobacterium tuberculosis

A pivotal moment occurred in 1882 when Robert Koch identified the causative agent of tuberculosis, *Mycobacterium tuberculosis*. This discovery laid the foundation for targeted diagnostic and therapeutic strategies, transforming tuberculosis from a mysterious disease into a scientifically understood condition.

Development of Diagnostic Techniques

Advancements such as the tuberculin skin test in the early 20th century allowed for better detection of latent and active infections. Improved microscopy and culture methods also enhanced the ability to diagnose and study the disease.

Introduction of Medical Treatments

Initial treatments, including sanatorium therapy emphasizing rest, nutrition, and fresh air, provided some relief but were insufficient for eradication. The discovery of streptomycin in 1943 marked the advent of antibiotic therapy, revolutionizing tuberculosis treatment.

Public Health Initiatives and Social Reforms

Sanatorium Movement and Isolation Policies

During the late 19th and early 20th centuries, sanatoria became central to TB management, emphasizing rest, nutrition, and fresh air. While beneficial, these facilities were often inaccessible to the impoverished, limiting their overall impact.

Improved Living Conditions

Urban reforms aimed at reducing overcrowding, improving sanitation, and promoting better nutrition contributed significantly to lowering transmission rates.

Vaccination Efforts

The development of the Bacillus Calmette-Guérin (BCG) vaccine in 1921 provided a preventive tool, especially effective in children, to reduce disease incidence.

The Impact of Antibiotics and Chemotherapy

Streptomycin and Subsequent Drugs

The discovery of streptomycin in 1943 was a breakthrough, followed by the development of other anti-tubercular drugs like isoniazid, rifampicin, and pyrazinamide. Combination therapy became the standard, significantly improving cure rates and reducing mortality.

Standardized Treatment Regimens

By the mid-20th century, standardized short-course chemotherapy regimens were implemented, leading to higher treatment success and decreased transmission.

Decline in Tuberculosis Incidence and Mortality

Statistical Trends (1850-1950)

Throughout this century, many countries experienced a sharp decline in TB cases and related deaths. For example, in the United States, tuberculosis mortality decreased from over 200 deaths per 100,000 population in the early 1900s to less than 20 per 100,000 by 1950.

Factors Contributing to the Retreat

- Effective antibiotic treatments
- Improved public health infrastructure
- Better socioeconomic conditions
- Widespread vaccination campaigns
- Enhanced diagnostic capabilities

Remaining Challenges and Ongoing Efforts

Persistent Reservoirs and Drug Resistance

Despite the successes, some regions continued to grapple with high TB burdens, compounded by issues such as multidrug-resistant TB (MDR-TB) and HIV co-infection.

Global Health Initiatives

Organizations like the World Health Organization launched comprehensive programs such as the DOTS (Directly Observed Treatment, Short-course) strategy to combat TB globally, emphasizing treatment adherence and surveillance.

Research and Future Directions

Ongoing research focuses on developing new vaccines, shorter treatment regimens, and molecular diagnostics to further reduce the disease's impact.

Conclusion

The period from 1850 to 1950 marked an extraordinary retreat of tuberculosis, driven by scientific discoveries, medical innovations, and public health reforms. The identification of *Mycobacterium tuberculosis*, the advent of antibiotics, and the implementation of vaccination and sanitation programs collectively transformed TB from a leading killer into a manageable disease in many parts of the world. While challenges remain, the lessons from this century-long fight continue to inform current strategies aimed at ending tuberculosis as a global health threat. The progress achieved during this era exemplifies how scientific progress combined with social action can significantly alter the course of infectious diseases, offering hope for future eradication efforts.

The retreat of tuberculosis 1850?1950 marks one of the most significant public health transformations in modern history. Over the course of a century, society witnessed a dramatic decline in the prevalence and mortality of tuberculosis (TB), transforming it from a leading cause of death to a largely controllable disease. This period saw a combination of scientific breakthroughs, public health initiatives, social changes, and medical advancements that collectively contributed to the retreat of TB. Analyzing this century-long journey provides insights into the progress of medicine, the importance of coordinated health policies, and the enduring challenges that remain in the fight against infectious diseases.

Introduction: The Historical Context of Tuberculosis (1850?1950)

In the 19th century, tuberculosis was often called the "white plague" due to its widespread devastation, especially in Europe and North America. It was a leading cause of death across all social classes, with no effective cure available. The industrial revolution, urbanization, and poor living conditions contributed to the rapid spread of the disease. By 1850, TB had become a major public health concern, prompting scientists, physicians, and governments to seek solutions.

The period from 1850 to 1950 was pivotal because it encompassed the dawn of microbiology, advances in medical science, and major shifts in public health policy. These developments collectively fostered a decline in TB cases and mortality, setting the foundation for modern disease control strategies.

Scientific Breakthroughs and the Understanding of Tuberculosis

The Discovery of the Mycobacterium tuberculosis

One of the most crucial milestones in the fight against TB was Robert Koch's discovery of Mycobacterium tuberculosis in 1882. This discovery was groundbreaking because:

- It confirmed the bacterial cause of TB, enabling targeted research.
- It shifted the perception of TB from a hereditary or environmental disease to an infectious one.
- It paved the way for diagnostic and vaccine development.

Features and Impact:

- Led to the development of bacteriological laboratories.
- Allowed for more accurate diagnosis.
- Sparked further research into bacterial physiology and pathogenicity.

Pros:

- Facilitated the development of specific diagnostic tools.
- Enabled epidemiological tracking.

Cons:

- Initially, it did not immediately translate into effective treatments.
- Bacterial identification required sophisticated laboratory infrastructure, limiting immediate global impact.

The Role of Microscopy and Diagnostics

Advances in microscopy, especially Ziehl-Neelsen staining (developed in 1883), allowed for easier identification of TB bacteria in clinical samples. This technological progress:

- Improved diagnostic accuracy.
- Allowed for early detection and isolation of infectious cases.
- Supported public health measures to contain outbreaks.

Features:

- Simple, cost-effective diagnostic method.
- Widely adopted in clinical settings.

Pros:

- Increased case detection.
- Facilitated epidemiological surveillance.

Cons:

- Limited sensitivity compared to modern molecular methods.
- Requires skilled technicians.

Development of Preventive Strategies

Vaccination: The Introduction of BCG

One of the most significant public health tools in reducing TB incidence was the development and deployment of vaccines. The Bacillus Calmette-Guérin (BCG) vaccine, introduced in 1921, represented a major step forward.

Features of BCG:

- Derived from an attenuated strain of *Mycobacterium bovis*.
- Administered primarily to infants and children.
- Provided protection against severe forms of childhood TB.

Pros:

- Safe and relatively inexpensive.
- Widely implemented in many countries.
- Reduced incidence of childhood TB and TB meningitis.

Cons:

- Variable efficacy against adult pulmonary TB.
- Not a sterilizing vaccine; vaccinated individuals could still develop disease.

Public Health Campaigns and Sanitation

Improvements in living conditions, sanitation, and nutrition played a pivotal role in the retreat of TB:

- Urban reforms reduced overcrowding and improving ventilation.
- Better nutrition boosted immune responses.
- Public health campaigns promoted awareness and early detection.

Features:

- Emphasis on social determinants of health.

- Implementation of TB clinics and mobile health units.

Pros:

- Addressed root causes of disease spread.
- Reduced transmission rates.

Cons:

- Required significant social policy changes.
- Progress was uneven and slow in some areas.

Medical Treatments and Their Impact

Pre-Antibiotic Era: Sanatoriums and Rest Therapy

Before antibiotics, treatment focused on rest, good nutrition, and fresh air. The sanatorium movement, prominent from the late 19th to early 20th centuries, aimed to isolate and treat TB patients in specialized facilities.

Features:

- Emphasis on prolonged rest and exposure to fresh air.
- Often located in rural or mountain settings.

Pros:

- Provided supportive care.
- Reduced transmission in some cases.

Cons:

- Limited efficacy; not curative.
- Expensive and inaccessible for many.
- Long hospital stays with no definitive cure.

The Antibiotic Revolution: Streptomycin and Beyond

The real turning point in TB control came with the discovery of antibiotics:

- Streptomycin (1943): The first effective antibiotic against TB.
- Followed by isoniazid, rifampicin, ethambutol, and pyrazinamide.

Features of Antibiotic Therapy:

- Multi-drug regimens to prevent resistance.
- Shorter treatment durations compared to sanatorium methods.
- Improved cure rates significantly.

Pros:

- Curative in most cases.
- Reduced infectiousness and transmission.
- Enabled outpatient treatment and community-based care.

Cons:

- Emergence of drug-resistant TB strains.
- Need for prolonged treatment adherence.
- Side effects and toxicity concerns.

The Decline of Tuberculosis: Quantitative and Qualitative Evidence

Statistical Trends and Impact

From 1850 to 1950, data from various countries illustrate a remarkable decline in TB incidence and mortality:

- Europe and North America: TB mortality decreased by up to 70-80% within this period.
- Epidemiological shifts: Disease shifted from being an almost universal childhood disease to more adult cases, owing to improved living conditions and vaccination.

Features:

- Declines correlated with urban reforms, vaccination, and antibiotics.
- Variability based on socioeconomic status and health policies.

Pros:

- Demonstrates the effectiveness of integrated public health measures.
- Validates scientific advances in disease control.

Cons:

- Data collection was inconsistent; some underreporting occurred.
- In some regions, progress was delayed due to social upheavals or resource limitations.

Remaining Challenges and Lessons Learned

Despite the tremendous progress, the retreat of TB was not absolute, and challenges persisted:

- Emergence of drug-resistant strains.
- Socioeconomic disparities affecting access to care.
- Need for better vaccines and shorter, more effective treatments.
- Ongoing transmission in vulnerable populations.

Lessons:

- Combating infectious diseases requires a multi-faceted approach combining science, social policy, and public engagement.
- Early detection and treatment are crucial.
- Investment in infrastructure and education yields long-term health benefits.

Conclusion: The Legacy of 1850-1950 in Tuberculosis Control

The period from 1850 to 1950 encapsulates a transformative era in the fight against tuberculosis. Scientific discoveries like Koch's identification of the causative bacterium and the advent of effective antibiotics fundamentally changed the disease landscape. Public health interventions,

improved living conditions, and vaccination campaigns contributed significantly to the decline in TB prevalence and mortality.

While the century witnessed remarkable progress, it also highlighted ongoing challenges, emphasizing that disease control is a continuous process. The lessons from this era laid the groundwork for modern tuberculosis control strategies, including the development of new vaccines, diagnostics, and treatments. The retreat of TB during this period exemplifies how coordinated scientific and social efforts can lead to profound improvements in public health, a legacy that continues to inform efforts against infectious diseases today.

Question What were the main factors contributing to the decline of tuberculosis between 1850 and 1950? The decline was primarily driven by improvements in public health measures, better sanitation, the development of effective sanitation infrastructure, and advances in medical treatment, including the discovery of antibiotics like streptomycin in the 1940s. How did the understanding of tuberculosis transmission evolve during the period 1850 to 1950? Initially misunderstood, the transmission was later understood to be primarily airborne, especially through inhaling droplets from coughs of infected individuals, which led to more targeted prevention strategies. What role did sanatoriums play in the retreat of tuberculosis during this period? Sanatoriums provided specialized treatment environments emphasizing rest, nutrition, and fresh air, which significantly contributed to controlling the disease before the advent of antibiotics. How did public health policies impact tuberculosis control from 1850 to 1950? Public health initiatives, including screening programs, isolation of infectious cases, and improved living conditions, helped reduce transmission and mortality rates over this century. What was the significance of the discovery of antibiotics in the decline of tuberculosis during this period? The discovery of antibiotics, particularly streptomycin in 1944, revolutionized treatment by effectively curing active TB cases, leading to a rapid decline in mortality and disease prevalence. Why is the period 1850-1950 considered a pivotal era in the history of tuberculosis control? It marks a transition from limited understanding and treatment to effective medical interventions and public health strategies, resulting in a significant retreat of the disease globally.

Related keywords: tuberculosis history, sanatorium movement, antimicrobial therapy, BCG vaccine, tuberculosis epidemiology, chest X-ray, tuberculin test, antibiotics, public health campaigns, tuberculosis eradication

Read the full article online: [The retreat of tuberculosis 1850 1950](#)

The ecology of mycobacteria impact on animal s an

The ecology of mycobacteria impact on animal health and ecosystems

Understanding the ecology of mycobacteria and their impact on animals is crucial for managing diseases, safeguarding biodiversity, and maintaining ecological balance. Mycobacteria are a diverse group of bacteria, some of which are pathogenic to animals and humans, while others exist benignly within various environmental reservoirs. Their complex interactions with animal hosts, environmental factors, and ecosystems influence disease transmission dynamics, wildlife health, and even public health. This article explores the ecological aspects of mycobacteria, their impact on animal populations, and the broader ecological implications.

Overview of Mycobacteria and Their Ecological Niche

What Are Mycobacteria?

Mycobacteria are a genus of bacteria characterized by their unique cell wall structure, which confers resistance to many disinfectants and environmental stresses. They include several notable pathogenic species such as *Mycobacterium tuberculosis*, *Mycobacterium bovis*, and *Mycobacterium avium*, as well as numerous non-pathogenic species. They are known for their ability to survive in diverse environments, including soil, water, and within host organisms.

Environmental Reservoirs and Survival Strategies

Mycobacteria thrive in various ecological niches, often persisting in environmental reservoirs that facilitate transmission to animal hosts. Key factors influencing their survival include:

- Resistance to desiccation and disinfectants due to lipid-rich cell walls
- Ability to form biofilms on surfaces and within soil or water matrices
- Persistence in soil, water, and organic matter for extended periods

These adaptations allow mycobacteria to maintain a stable ecological presence, making them persistent sources of infection.

Impact of Mycobacteria on Animal Populations

Pathogenicity and Disease Dynamics

Several mycobacterial species are pathogenic to animals, causing diseases that range from chronic wasting to severe systemic infections. The impact on animal populations can be significant, especially in wildlife and livestock.

- **Bovine tuberculosis (*Mycobacterium bovis*):**

- Primarily affects cattle, deer, and wildlife such as badgers and badgers
- Leads to weight loss, cough, and decreased productivity
- Can transmit to humans through unpasteurized milk or close contact

- ***Mycobacterium avium* complex (MAC):**

- Impacts poultry, pigs, and other domestic animals
- Causes chronic infections, especially in immunocompromised animals

- **Wildlife and non-tuberculous mycobacteria:**

- Can cause skin lesions, lymphadenitis, and systemic illness
- May lead to population declines in vulnerable species

Transmission Pathways and Ecology

Mycobacteria spread among animals through various pathways, often influenced by environmental factors:

- Ingestion of contaminated water or feed
- Direct contact with infected animals or contaminated environments
- Inhalation of aerosolized bacteria in enclosed spaces or dense populations

Environmental reservoirs such as contaminated soil and water bodies serve as continuous sources, especially in areas with high animal density.

Ecological Factors Influencing Mycobacterial Dynamics

Environmental Conditions

The stability and proliferation of mycobacteria are heavily influenced by ecological factors:

- **Temperature:** Moderate temperatures promote persistence; extreme heat or cold can reduce viability
- **Moisture:** Adequate moisture levels support bacterial survival in soil and water
- **pH levels:** Slightly acidic to neutral pH often favors survival, while extreme pH levels inhibit

growth

Ecological Interactions

Interactions with other microorganisms and environmental components shape the ecology of mycobacteria:

- Biofilm formation enhances resistance and persistence
- Competition with other soil microbes can limit or promote mycobacterial survival
- Presence of organic matter provides nutrients, supporting growth

Broader Ecological and Conservation Impacts

Wildlife Health and Biodiversity

Mycobacterial infections can threaten wildlife populations, especially species with limited ranges or those already under ecological stress:

- Chronic infections reduce fitness and reproductive success
- Outbreaks may lead to population declines or shifts in community dynamics
- Infections can serve as indicators of ecosystem health deterioration

Transmission Between Wildlife, Livestock, and Humans

The ecology of mycobacteria facilitates cross-species transmission, posing challenges for disease control:

- Wildlife reservoirs can infect livestock, complicating eradication efforts
- Contaminated environments serve as common sources for multiple species
- Human activities, such as farming and hunting, increase exposure risks

Implications for Ecosystem Management

Effective management of mycobacterial diseases requires understanding their ecological context:

- Monitoring environmental reservoirs and wildlife health
- Implementing biosecurity measures to reduce environmental contamination

- Conservation strategies to protect vulnerable species from infection

Strategies for Managing Mycobacterial Impact in Ecosystems

Environmental Control Measures

To mitigate the ecological impact of mycobacteria, targeted strategies include:

- Reducing environmental contamination through proper waste management
- Controlling wildlife-livestock interactions in high-risk areas
- Restoring natural habitats to support resilient ecosystems

Surveillance and Research

Ongoing research enhances understanding of mycobacterial ecology:

- Mapping environmental reservoirs and hotspots
- Studying transmission pathways in different ecological settings
- Developing sensitive diagnostic tools for environmental and biological samples

One Health Approach

Addressing the ecological impact of mycobacteria requires integrated efforts:

- Coordination among veterinary, environmental, and public health sectors
- Community engagement and education on disease prevention
- Policy development for sustainable ecosystem management

Conclusion

The ecology of mycobacteria significantly influences their impact on animal health and ecosystems. Their ability to persist in environmental reservoirs, adapt to diverse ecological niches, and transmit across species underscores the importance of understanding their ecological dynamics. Effective management and control strategies hinge on a comprehensive appreciation of these bacteria's ecological roles, highlighting the need for multidisciplinary approaches that integrate environmental health, wildlife conservation, livestock management, and public health. By advancing our knowledge in this field, we can better protect animal populations, preserve biodiversity, and reduce the risks of zoonotic transmission, ensuring healthier ecosystems for future generations.

The ecology of mycobacteria impact on animal populations is a topic of increasing scientific interest, given the significant implications for wildlife health, livestock management, and zoonotic disease transmission. Mycobacteria, a diverse genus of bacteria known for their hardy nature and capacity to cause chronic infections, play a complex role within various ecological niches. Understanding their ecology?the interactions among mycobacteria, animal hosts, environmental factors, and human activity?is crucial for developing effective control strategies and mitigating their impact on animal populations.

Introduction to Mycobacteria and Their Ecological Significance

Mycobacteria encompass over 200 species, including notable pathogens such as *Mycobacterium tuberculosis*, responsible for tuberculosis (TB) in humans, and *Mycobacterium bovis*, which causes bovine TB in cattle and can infect multiple species. Some species are environmental saprophytes, thriving in soil and water, while others are obligate pathogens. Their ecological significance extends beyond direct pathogenicity, influencing biodiversity, ecosystem health, and the dynamics of infectious diseases in animal populations.

Understanding the ecology of mycobacteria involves examining their survival mechanisms, transmission pathways, environmental reservoirs, and interactions with hosts across various ecosystems. These elements collectively determine how mycobacteria persist, spread, and impact animal health at local and global scales.

Environmental Reservoirs and Survival Strategies of Mycobacteria

Environmental Reservoirs

Many mycobacteria, especially the non-tuberculous mycobacteria (NTM), are environmental organisms found in soil, water, and sediment. These reservoirs serve as a continual source of exposure for animals and humans. For example:

- Soil: Rich in organic matter, soil provides a habitat for NTM such as *Mycobacterium avium* complex (MAC).
- Water bodies: Lakes, ponds, and wetlands can harbor pathogenic mycobacteria, especially in biofilms.
- Fomites and fomite surfaces: Contaminated feed, bedding, and equipment can act as vectors.

Environmental persistence is facilitated by the unique cell wall structure of mycobacteria, which contains mycolic acids that confer resistance to desiccation, disinfectants, and varying temperature conditions.

Survival Mechanisms

Mycobacteria have evolved several strategies to survive in hostile environments:

- Biofilm formation: Aggregates embedded in a self-produced matrix enhance resistance to environmental stressors and immune responses.
- Dormancy and low metabolic activity: Some species can enter a dormant state, enabling long-term survival under unfavorable conditions.
- Resistance to disinfectants: Their cell wall's lipid-rich composition makes them inherently resistant to many common disinfectants.

These adaptations allow mycobacteria to persist in environmental niches for extended periods, increasing the likelihood of contact with susceptible animal hosts.

Transmission Dynamics and Pathways in Animal Populations

Infection Routes

Mycobacteria can infect animals through multiple pathways:

- Ingestion: Consuming contaminated water or feed contaminated with mycobacteria is a primary route, especially for NTM.
- Inhalation: Aerosolized bacteria from environmental sources or infected animals can infect respiratory tissues.
- Trauma and skin contact: Open wounds or skin abrasions can serve as entry points.

The transmission efficiency varies across species and depends on environmental factors and host susceptibility.

Host Susceptibility and Disease Progression

Animal susceptibility to mycobacterial infections depends on:

- Species and breed: Some species are more resistant or susceptible; for instance, cattle are highly susceptible to *M. bovis*, whereas some wildlife species may act as reservoirs.
- Age: Younger animals often exhibit higher susceptibility.
- Immune status: Stress, co-infections, and nutritional deficiencies can impair immune responses, facilitating infection.

Once infected, animals often develop a latent or active infection, with clinical signs ranging from subclinical to severe, chronic disease.

Impact of Mycobacteria on Animal Ecology

Effects on Wildlife Populations

Wildlife species can serve as reservoirs or spillover hosts for pathogenic mycobacteria, influencing disease dynamics:

- Reservoir hosts: Species such as badgers, deer, and bison have been identified as maintaining *M. bovis* populations, complicating eradication efforts in livestock.
- Disease spillover: Infection in wildlife can spill over to domestic animals, creating a cycle of transmission that is difficult to control.
- Population health: Chronic mycobacterial infections can reduce fitness, reproductive success, and survival, impacting population stability.

The presence of mycobacteria can also alter predator-prey dynamics, with infected animals being more vulnerable or less competitive.

Impact on Livestock and Domestic Animals

In farm settings, mycobacterial infections have multifaceted impacts:

- Economic losses: Reduced productivity, culling, and trade restrictions.
- Animal welfare: Chronic disease leads to suffering and decreased quality of life.
- Management challenges: Difficulties in eradication due to environmental reservoirs and wildlife reservoirs.

The ecology of mycobacteria influences disease persistence and complicates control measures in agricultural contexts.

Environmental and Anthropogenic Factors Influencing Mycobacteria Ecology

Climate and Geographic Variability

Temperature, humidity, and rainfall patterns affect the survival and distribution of mycobacteria:

- Warm, humid environments favor biofilm formation and bacterial persistence.
- Arid conditions may reduce environmental loads but can concentrate bacteria in water sources or sediments.

Geographic features influence the interaction between wildlife, livestock, and environmental reservoirs, shaping regional disease dynamics.

Human Activities and Land Use

Human interventions significantly impact mycobacterial ecology:

- Agricultural practices: Overstocking, poor sanitation, and feed management can increase exposure.
- Wildlife management: Encroachment, habitat fragmentation, and pollution can alter wildlife-livestock interfaces.
- Water management: Use of contaminated water sources can facilitate transmission.

Urbanization and deforestation can disturb ecological balances, leading to increased contact zones and transmission opportunities.

Control Strategies and Ecological Considerations

Vaccination and Diagnostic Tools

Understanding the ecology of mycobacteria informs control efforts:

- Vaccination: Bovine TB control programs rely on vaccines, but environmental persistence can limit effectiveness.
- Diagnostics: Accurate detection of latent and active infections is essential; environmental sampling can help identify reservoirs.

Environmental Management and Biosecurity

Strategies include:

- Reducing environmental contamination: Proper disposal of carcasses, sanitation of equipment, and water treatment.
- Habitat management: Limiting wildlife-livestock contact through fencing and habitat modification.

- Monitoring and surveillance: Ecological surveillance provides insights into hotspots and transmission pathways.

Integrative Approaches

Effective management requires a One Health approach, integrating veterinary, ecological, and public health perspectives. Ecological studies can help predict disease emergence and guide targeted interventions.

Future Directions and Research Needs

Advancements in ecological modeling, genomics, and environmental microbiology are expanding our understanding of mycobacterial ecology. Key areas for future research include:

- Mapping environmental reservoirs across different ecosystems.
- Understanding host-pathogen-environment interactions that facilitate persistence and spread.
- Developing eco-friendly control measures that consider ecological balances.
- Assessing the impact of climate change on the distribution and ecology of mycobacteria.

Such insights will enhance our capacity to mitigate their impact on animal health and prevent zoonotic spillovers.

Conclusion

The ecology of mycobacteria profoundly influences their persistence, transmission, and impact on animal populations. Their ability to survive in diverse environmental reservoirs, adapt to various hosts, and respond to ecological and anthropogenic changes makes them formidable pathogens within the animal health landscape. A comprehensive understanding of their ecological dynamics is essential for designing effective control measures, safeguarding biodiversity, and preventing zoonotic transmission. As research advances, integrating ecological insights with veterinary and public health efforts will be paramount in managing mycobacterial diseases in a rapidly changing world.

Question What role do mycobacteria play in the ecology of animal populations? Mycobacteria influence animal populations by causing diseases such as tuberculosis and leprosy, which can affect individual health, population dynamics, and interactions within ecosystems. How do environmental factors affect the persistence of mycobacteria in animal habitats? Environmental factors like soil composition, humidity, temperature, and water sources can promote or inhibit the survival of mycobacteria, impacting their transmission among animals. What is the significance of wildlife reservoirs in the ecology of mycobacteria? Wildlife reservoirs serve as sources of

mycobacterial infection for domestic animals and humans, maintaining the bacteria in nature and complicating control efforts. How do mycobacteria impact the health and behavior of affected animals? Infected animals may experience weight loss, weakness, and respiratory issues, which can alter their behavior, reduce reproductive success, and increase mortality rates. What environmental transmission pathways facilitate the spread of mycobacteria among animals? Transmission pathways include contaminated water, soil, feed, and direct contact with infected animals or their excreta, enabling mycobacteria to persist and spread in various environments. How does the ecology of mycobacteria influence disease management in animal populations? Understanding their ecological niches helps in designing targeted control strategies, such as habitat management, vaccination, and biosecurity measures to reduce transmission. Are there specific ecological traits of mycobacteria that enhance their survival in natural environments? Yes, mycobacteria's hardy cell wall, ability to form biofilms, and resistance to environmental stresses contribute to their persistence outside hosts. What is the impact of mycobacterial infections on biodiversity and ecosystem health? Mycobacterial diseases can reduce animal diversity by causing mortality in susceptible species, potentially disrupting ecological balances and impacting ecosystem services. How can studying the ecology of mycobacteria inform better control and prevention strategies in animals? Ecological studies reveal transmission dynamics and environmental reservoirs, guiding interventions like habitat modification, vaccination programs, and monitoring to prevent outbreaks.

Related keywords: mycobacteria, ecology, animal health, zoonotic diseases, tuberculosis, environmental reservoirs, pathogen transmission, wildlife, infectious diseases, disease ecology

Read the full article online: [The Ecology Of Mycobacteria Impact On Animal S An](#)

tropical surgery vademecum a vademacum series boo

tropical surgery vademecum a vademacum series boo is an essential resource for healthcare professionals working in tropical medicine and surgery. As tropical regions often present unique challenges due to endemic diseases, environmental factors, and resource limitations, having a comprehensive, well-structured guide like the Vademecum Series Boo becomes invaluable. This article provides an in-depth exploration of the Tropical Surgery Vademecum, its significance, key features, and practical applications for medical practitioners aiming to improve patient outcomes in tropical settings.

Understanding the Tropical Surgery Vademecum Series Boo

What is the Vademecum Series Boo?

The Vademecum Series Boo is a specialized collection of medical reference books designed for quick consultation and practical guidance across various medical disciplines. The "Tropical Surgery" volume focuses on surgical procedures, conditions, and management strategies pertinent to tropical environments. It synthesizes current research, clinical best practices, and resource-efficient techniques tailored for settings with limited infrastructure.

Purpose and Audience

The primary aim of the Tropical Surgery Vademecum is to serve as an indispensable tool for:

- Surgeons working in tropical regions
- General practitioners handling surgical emergencies
- Medical students and residents specializing in tropical medicine
- Healthcare policymakers designing intervention programs

The guide emphasizes user-friendly language, clear illustrations, and concise protocols to facilitate swift decision-making.

Key Features of the Tropical Surgery Vademecum

Comprehensive Coverage

The Vademecum covers a wide array of topics, including:

- Traumatic injuries common in tropical zones
- Infectious disease-related surgical conditions
- Obstetric and gynecological surgeries
- Pediatric surgical issues unique to tropical settings
- Management of tropical-specific emergencies such as snakebites and envenomations

Practical and Resource-Conscious Approaches

Given the resource limitations often encountered in tropical healthcare facilities, the guide:

- Recommends alternative techniques when modern equipment is unavailable
- Emphasizes the use of locally available materials
- Prioritizes cost-effective interventions without compromising patient safety

Illustrations and Step-by-Step Protocols

Clear diagrams, flowcharts, and checklists are integrated throughout the manual to:

- Facilitate understanding of complex procedures
- Ensure standardization of care
- Reduce errors during emergency situations

Updated and Evidence-Based Content

The latest edition incorporates recent research findings, including:

- Advances in surgical techniques
- Emerging tropical diseases impacting surgical practice
- Updated antibiotic and antimicrobial guidelines

Core Topics Covered in the Vademecum

Trauma and Emergency Surgery

Tropical regions often see higher incidences of trauma due to:

- Road traffic accidents
- Animal bites
- Natural disasters

The Vademecum offers protocols for:

- Wound management
- Fracture stabilization
- Triage and emergency surgical interventions

Infectious Disease-Related Surgical Conditions

Infections endemic to tropical areas frequently complicate surgical care:

- Mycetoma and sporotrichosis
- Hydatid cysts
- Buruli ulcer
- Leprosy-related deformities

The guide provides insights on:

- Diagnosis
- Surgical excision techniques
- Postoperative care

Obstetric and Gynecological Surgeries

Complications such as obstructed labor, postpartum hemorrhage, and endemic infections require specialized surgical management:

- Cesarean sections
- Hysterectomy procedures
- Management of obstetric fistulas

Pediatric Surgical Conditions

Common pediatric issues include:

- Congenital anomalies
- Infectious abscesses
- Trauma-related injuries

The manual emphasizes age-appropriate anesthesia and postoperative care strategies.

Tropical-Specific Emergencies

Critical conditions requiring immediate surgical intervention include:

- Snakebite envenomation with tissue necrosis
- Malaria-associated splenic rupture
- Filariasis-related lymphatic obstructions

Application of the Vademecum in Clinical Practice

Preoperative Planning

The guide assists clinicians in:

- Assessing the severity of tropical diseases
- Selecting appropriate surgical techniques
- Preparing resource-efficient surgical plans

Intraoperative Management

Step-by-step instructions help ensure:

- Sterile technique adherence
- Efficient use of available equipment
- Management of intraoperative complications

Postoperative Care and Follow-up

Recommendations include:

- Infection control measures tailored for tropical pathogens
- Wound care protocols
- Rehabilitation strategies in resource-limited settings

Benefits of Using the Tropical Surgery Vademecum

- **Improved Patient Outcomes:** By providing evidence-based, practical guidance, the manual enhances surgical success rates and reduces complications.
- **Enhanced Decision-Making:** Clear protocols enable clinicians to make swift, informed choices, especially during emergencies.
- **Resource Optimization:** Emphasizing low-cost, locally available solutions makes high-quality care accessible even in under-resourced facilities.
- **Educational Value:** The manual serves as a teaching tool for students and trainees, fostering consistent surgical practices.

Integrating the Vademecum into Healthcare Systems

Training and Capacity Building

Healthcare providers can utilize the Vademecum for:

- Workshops and training sessions
- Continuing medical education
- Developing standardized surgical protocols

Policy and Program Development

Health authorities can reference the manual to:

- Design surgical intervention programs tailored to tropical diseases
- Allocate resources efficiently
- Establish quality assurance standards

Conclusion: Why the Tropical Surgery Vademecum Series Boo Is Indispensable

The Tropical Surgery Vademecum in the Vademecum Series Boo stands out as a crucial resource for addressing the unique surgical challenges posed by tropical environments. Its comprehensive content, practical guidance, and resource-conscious approach make it an essential tool for clinicians committed to improving surgical care in regions where resources are limited, and disease burden is high.

By integrating the insights and protocols from this manual, healthcare practitioners can enhance their clinical practice, reduce surgical morbidity and mortality, and ultimately contribute to better health outcomes in tropical populations. Whether used as a quick reference during emergencies or as a teaching aid for training future surgeons, the Vademecum Series Boo remains a cornerstone of tropical surgical medicine.

Keywords: Tropical Surgery, Vademecum Series Boo, tropical medicine, surgical guide, resource-limited settings, tropical diseases, emergency surgery, surgical protocols, tropical healthcare, medical reference.

Tropical Surgery Vademecum: A Vademecum Series Book

In the landscape of global health, tropical medicine and surgery stand as a critical frontier, addressing complex surgical challenges in resource-limited, tropical environments. The Tropical Surgery Vademecum, part of the esteemed Vademecum Series, emerges as a comprehensive guide designed for practitioners, students, and healthcare providers operating in these demanding settings. This publication encapsulates practical knowledge, surgical techniques, and contextual insights tailored specifically to the unique needs of tropical regions.

Introduction to the Vademecum Series and Its Significance

The Vademecum Series is renowned for its pragmatic approach to medical knowledge, offering concise yet thorough references across various specialties. The Tropical Surgery Vademecum continues this tradition, focusing exclusively on surgical issues prevalent in tropical zones?areas often characterized by endemic diseases, limited infrastructure, and diverse pathologies.

Why a Specialized Surgical Guide for Tropical Regions?

Traditional surgical textbooks often overlook the nuances of tropical environments. These regions pose unique challenges such as:

- High prevalence of infectious diseases: Malaria, leishmaniasis, filariasis, and other parasitic infections often complicate surgical interventions.
- Resource constraints: Limited availability of advanced imaging, sterile supplies, and specialized equipment.
- Environmental factors: High humidity, temperature fluctuations, and endemic pests influence wound healing and infection control.
- Cultural considerations: Variations in health beliefs and practices impact patient management and compliance.

The Tropical Surgery Vademecum addresses these factors, providing tailored strategies to optimize surgical outcomes under such conditions.

Core Content and Structure of the Vademecum

The book is organized into sections that systematically cover essential aspects of tropical surgery, from preoperative assessment to postoperative care, integrating disease-specific chapters.

- Preoperative Evaluation in Tropical Settings

Effective surgical management begins with thorough evaluation, adapted to the realities of tropical

regions.

- History Taking and Clinical Examination: Emphasis on endemic disease exposure, travel history, and occupational risks.
- Laboratory and Diagnostic Limitations: Guidance on clinical diagnosis when advanced labs are unavailable.
- Risk Stratification: Identifying patients with comorbidities such as malnutrition, anemia, or infections that influence surgical risk.

- Surgical Techniques Adapted for Resource-Limited Environments

The guide provides detailed descriptions of procedures with modifications suitable for limited-resource settings.

- Wound Management: Techniques for managing tropical ulcers, mycetomas, and other infectious wounds.
 - Trauma Surgery: Strategies for managing injuries common in tropical regions, including animal bites and blast injuries.
 - Lymphatic and Vascular Surgery: Addressing filarial lymphedema and vascular anomalies.
- #### - Disease-Specific Surgical Approaches

The core of the book is dedicated to prevalent diseases requiring surgical intervention:

- Buruli Ulcer: Surgical excision, skin grafting, and rehabilitation.
 - Leishmaniasis: Nodule excision and management of secondary infections.
 - Filarial Lymphedema: Debulking procedures and limb reconstruction.
 - Trauma and Emergency Surgery: Handling of snake bites, animal bites, and other injuries.
- #### - Postoperative Care and Follow-up

Post-surgical management is tailored to tropical environments:

- Infection Control: Use of locally available antiseptics, wound dressings, and antibiotics.

- Nutritional Support: Addressing malnutrition to promote healing.
- Rehabilitation: Techniques to prevent disability, especially in limb surgeries.
- Preventive and Public Health Aspects

Beyond individual patient care, the guide emphasizes preventive measures:

- Community education on early presentation and wound care.
- Vector control strategies to reduce disease transmission.
- Vaccination and prophylactic measures where applicable.

Key Features of the Vademecum

The Tropical Surgery Vademecum stands out for its practical orientation and user-friendly design:

- Concise Summaries: Clear, bullet-pointed instructions for rapid reference.
- Photographs and Illustrations: Visual aids illustrating surgical techniques, wound management, and disease presentations.
- Checklists and Algorithms: Step-by-step guides for decision-making processes.
- Case Studies: Real-world scenarios illustrating common challenges and solutions.

Importance for Global Health and Surgical Training

The book serves as a vital resource for surgeons working in tropical regions, whether in Africa, Southeast Asia, Latin America, or island nations. It bridges the gap between ideal surgical practice and the realities faced on the ground.

For Surgeons and Healthcare Workers

- Enhances preparedness for managing tropical surgical pathologies.
- Provides guidance on improvisation and resourcefulness.
- Supports training programs by offering standardized protocols.

For Researchers and Public Health Officials

- Offers insights into prevalent surgical diseases and their management.

- Guides the development of health policies and surgical outreach programs.
- Encourages integration of surgical care into broader tropical disease control strategies.

Challenges Addressed by the Vademecum

The Tropical Surgery Vademecum confronts critical issues faced in these settings:

- Limited Resources: Techniques requiring minimal equipment.
- High Disease Burden: Focused management of endemic pathologies.
- Infection Control: Strategies adapted to high infection risks.
- Training Gaps: Aimed at equipping non-specialist surgeons and general practitioners.

By emphasizing adaptability and practicality, the guide empowers healthcare providers to deliver effective surgical care despite constraints.

Future Directions and Continuing Relevance

As tropical regions evolve and new challenges emerge?such as emerging infectious diseases or climate change impacts?the Vademecum Series continues to adapt. The Tropical Surgery Vademecum is expected to incorporate updates on minimally invasive techniques, telemedicine integration, and new disease patterns.

Its role remains pivotal in education, capacity building, and improving surgical outcomes in underserved tropical communities.

Conclusion

The Tropical Surgery Vademecum from the Vademecum Series exemplifies a dedicated effort to equip healthcare providers with the knowledge and tools necessary for effective surgical intervention in challenging tropical environments. Its comprehensive, practical approach ensures that even in resource-limited settings, patients receive quality surgical care tailored to their unique epidemiological and environmental context. As global health continues to prioritize equitable access and sustainable practices, such specialized guides will remain indispensable resources for advancing surgical care in the tropics.

Question What is the 'Tropical Surgery Vademecum' and who is it intended for? The 'Tropical Surgery Vademecum' is a comprehensive guide designed for healthcare professionals working in tropical regions, providing essential surgical protocols and management strategies tailored to resource-limited settings. **Answer** How does the 'Vademecum Series Boo' enhance surgical practice in tropical environments? The 'Vademecum Series Boo' offers concise, evidence-based

protocols, quick reference charts, and practical tips that streamline surgical decision-making and improve patient outcomes in tropical healthcare settings. What are the key topics covered in the 'Tropical Surgery Vademecum'? The manual covers topics such as infectious disease management, trauma care, surgical techniques for endemic conditions, anesthesia considerations, and postoperative care specific to tropical regions. Is the 'Tropical Surgery Vademecum' suitable for both students and experienced surgeons? Yes, the vademecum is designed to be accessible for students, residents, and experienced surgeons, providing foundational knowledge and advanced guidance tailored to tropical surgical challenges. Are there digital versions or mobile applications for the 'Vademecum Series Boo'? Yes, the series is often available in digital formats and mobile apps, allowing healthcare providers to access vital information quickly in remote or resource-limited settings. How frequently is the 'Tropical Surgery Vademecum' updated to reflect new guidelines? The manual is updated periodically to incorporate the latest evidence-based practices, emerging diseases, and regional health considerations, ensuring clinicians have current information. Does the 'Tropical Surgery Vademecum' include case studies or real-world scenarios? Yes, it features case studies and scenario-based guidance that help practitioners apply theoretical knowledge to practical situations commonly encountered in tropical surgical practice. Can the 'Vademecum Series Boo' be used as a training resource in tropical surgery workshops? Absolutely, it is an excellent resource for training sessions, workshops, and continuing medical education programs focused on tropical surgical care. Where can healthcare professionals obtain the 'Tropical Surgery Vademecum' and the 'Vademecum Series Boo'? These resources are typically available through medical publishers, specialized tropical medicine organizations, and online platforms dedicated to global health and surgical education.

Related keywords: tropical surgery, medical vademecum, surgical manual, tropical medicine, clinical guide, medical reference, vademecum series, surgical handbook, tropical health, medical textbook

Read the full article online: [Tropical Surgery Vademecum A Vademecum Series Boo](#)

forgotten people forgotten diseases the neglected

forgotten people forgotten diseases the neglected: Shedding Light on the Overlooked and Underserved

In a world where medical advancements and global health initiatives often focus on high-profile diseases like COVID-19, cancer, and HIV/AIDS, a crucial segment of health challenges remains largely invisible: the forgotten people and forgotten diseases. These are the marginalized populations suffering from neglected diseases that rarely make headlines, receive insufficient

funding, and often lack adequate treatment options. Addressing these issues is vital for achieving health equity and ensuring that no one is left behind in our collective quest for well-being. This article explores the concept of neglected diseases, highlights the vulnerable populations affected, and underscores the importance of global efforts to combat these overlooked health threats.

Understanding Neglected Diseases and Their Impact

Neglected diseases are a group of infectious diseases that predominantly affect impoverished communities in developing countries. Unlike well-known illnesses that attract significant research and funding, neglected diseases often remain in the shadows, perpetuating cycles of poverty and ill health.

What Are Neglected Diseases?

Neglected diseases include a wide range of infections caused by bacteria, parasites, viruses, and fungi. They are characterized by:

- Limited research and development efforts
- Minimal attention from pharmaceutical companies due to limited profitability
- High prevalence in underserved populations
- Significant social and economic burdens

Some of the most common neglected diseases are:

- Chagas disease
- Leishmaniasis
- Human African trypanosomiasis (sleeping sickness)
- Schistosomiasis
- Lymphatic filariasis (elephantiasis)
- Onchocerciasis (river blindness)
- Buruli ulcer
- Leprosy

The Global Burden of Neglected Diseases

Despite their name, neglected diseases impose a substantial health burden worldwide. The World Health Organization estimates that over one billion people are affected by at least one neglected tropical disease (NTD). These diseases often cause chronic disability, disfigurement, and sometimes death, creating a cycle of poverty for affected individuals and their communities.

The economic impacts are equally devastating. Treatment costs, lost productivity, and the social stigma associated with these diseases hinder community development and exacerbate inequality.

Forgotten People: The Marginalized and Vulnerable

Neglected diseases primarily afflict people living in extreme poverty, often in rural or marginalized urban settings. These populations face multiple barriers to healthcare access, including lack of infrastructure, insufficient health education, and cultural stigmas.

Who Are the Forgotten People?

The key groups most affected include:

- **Residents of impoverished rural communities:** Limited access to healthcare services and clean water increases exposure to disease vectors.
- **Indigenous populations:** Often marginalized, with less access to health systems and higher vulnerability to neglected diseases.
- **Refugees and displaced persons:** Crowded living conditions and poor sanitation contribute to disease spread.
- **Children and women:** Particularly susceptible to the long-term impacts of neglected diseases such as lymphatic filariasis and leprosy.

These populations often remain invisible to mainstream health initiatives, which focus more on high-profile health challenges.

Social and Economic Consequences

Neglected diseases can cause disfigurement, disabilities, and chronic health issues that hinder education, employment, and social participation. The social stigma associated with visible symptoms like disfigurement or disability further isolates affected individuals, perpetuating cycles of poverty and marginalization.

Furthermore, the lack of awareness and resources means that many people are unaware of prevention methods or available treatments, worsening health outcomes.

Challenges in Addressing Forgotten Diseases

Despite the significant burden, several barriers hinder effective management and eradication of neglected diseases.

Limited Funding and Research

Pharmaceutical companies often lack incentives to develop vaccines or treatments for diseases that predominantly affect impoverished populations. Consequently, research and development (R&D) are underfunded, leading to a scarcity of effective medicines and diagnostics.

Weak Healthcare Infrastructure

Many affected regions lack basic healthcare infrastructure, including clinics, trained health workers, and supply chains for medicines. This impedes timely diagnosis and treatment.

Stigma and Cultural Barriers

Cultural beliefs and social stigma can discourage individuals from seeking medical help, especially for diseases that cause visible disfigurement or are associated with shame.

Inadequate Policy and Global Attention

Neglected diseases often do not receive the political will or funding necessary for large-scale intervention programs, partly due to competing priorities and limited awareness.

Global Efforts and Solutions

While challenges abound, concerted global efforts are making progress in addressing neglected diseases and their affected populations.

International Initiatives and Campaigns

Organizations like the World Health Organization (WHO), the Bill & Melinda Gates Foundation, and various NGOs have launched initiatives targeting neglected diseases with strategies including:

- Development of new diagnostics, medicines, and vaccines
- Mass drug administration programs
- Community education and engagement
- Strengthening healthcare infrastructure

The London Declaration on Neglected Tropical Diseases (2012), for example, committed global partners to control or eliminate several NTDs by 2020.

Innovations and Research

Advances in technology, such as mobile health platforms and genomic research, are improving disease detection and treatment options. Additionally, pharmaceutical companies are increasingly investing in neglected disease R&D through public-private partnerships.

Community-Centered Approaches

Empowering local communities through education, participation, and culturally sensitive interventions enhances the effectiveness of disease control measures.

How You Can Help

Addressing these overlooked issues requires collective effort. Here are ways individuals and organizations can contribute:

- **Raise awareness:** Educate others about neglected diseases and their impact.
- **Support NGOs:** Donate to organizations working in affected regions.
- **Advocate for policy change:** Push for increased funding and attention to neglected diseases.
- **Volunteer:** Offer skills or resources to health initiatives in underserved areas.

Every action, big or small, can contribute to breaking the cycle of neglect and suffering.

Conclusion

Forgotten people, forgotten diseases, the neglected ? these words encapsulate a vital aspect of global health that demands more attention, compassion, and action. The populations affected by neglected diseases are among the most vulnerable, often invisible in mainstream health narratives. Their struggles highlight the urgent need for increased research, better healthcare infrastructure, and sustained international commitment. By shining a spotlight on these overlooked issues, we can foster a more equitable world where health is a universal right, not a privilege reserved for the few. Together, through awareness, advocacy, and innovation, we can ensure that no one remains forgotten in the fight against disease.

Forgotten People, Forgotten Diseases: The Neglected Crisis

In the vast landscape of global health, there exists a shadowy realm often overlooked?neglected diseases and the communities most affected by them. These illnesses, despite their profound impact, remain underfunded, under-researched, and under-prioritized on the world stage. The term ?neglected? extends beyond the diseases themselves to the populations suffering from them?many of whom are marginalized, impoverished, or living in remote areas. This comprehensive exploration aims to shed light on these overlooked ailments and the people they afflict, emphasizing the urgent

need for awareness, research, and intervention.

Understanding the Scope of Neglected Diseases

What Are Neglected Diseases?

Neglected diseases, often called Neglected Tropical Diseases (NTDs), are a diverse group of communicable diseases that predominantly affect impoverished populations in developing regions. They are characterized by:

- High prevalence in impoverished communities: Often linked to poor sanitation, limited access to healthcare, and environmental factors.
- Low research and drug development investment: They receive a fraction of the funding allocated to more prominent diseases like HIV/AIDS, malaria, or tuberculosis.
- Chronic and debilitating nature: Many cause long-term health issues, disability, and social stigma.
- Limited awareness: They are rarely discussed in mainstream media or global health agendas.

The World Health Organization (WHO) recognizes over 20 diseases classified as NTDs, including:

- Chagas disease
- Leishmaniasis
- Schistosomiasis
- Lymphatic filariasis
- Onchocerciasis (river blindness)
- Soil-transmitted helminth infections
- Trachoma
- Dengue and chikungunya (though more prominent, still underfunded)

Beyond NTDs, other neglected diseases include leprosy, rabies, and certain parasitic infections, all contributing to a cycle of poverty and disease.

The People Left Behind: Communities Suffering from Neglected Diseases

Who Are the Forgotten People?

The populations most affected by neglected diseases are often marginalized for various reasons:

- Poverty and Socioeconomic Disadvantage: Limited access to healthcare, education, and sanitation.
- Rural and Remote Locations: Communities living in inaccessible regions with minimal healthcare infrastructure.
- Indigenous and Ethnic Minorities: Often face discrimination, cultural barriers, and neglect.
- Children and Women: Particularly vulnerable due to biological, social, and economic factors.
- Laborers and Agricultural Workers: Exposure to disease vectors in work environments.

Many of these groups are invisible in national and global health reports, making targeted intervention challenging. Their struggles are compounded by systemic neglect, leading to a persistent cycle of disease and poverty.

Case Studies of Neglected Communities

- The Amazon Basin Indigenous Communities: Suffer from Chagas disease and leishmaniasis due to vector exposure and limited healthcare.
- Sub-Saharan African Rural Populations: Endemic for schistosomiasis and lymphatic filariasis, with minimal access to treatment.
- South Asian Slum Dwellers: Often face soil-transmitted helminth infections and dengue outbreaks.
- Refugee Camps and Displacement Sites: High risk for multiple NTDs due to overcrowding and poor sanitation.

Deep Dive into Specific Neglected Diseases

Chagas Disease

Overview: Caused by the parasite *Trypanosoma cruzi*, transmitted mainly through the triatomine bug (?kissing bug?), Chagas disease affects an estimated 6-7 million people globally, primarily in Latin America.

Impact on People:

- Chronic cardiac and digestive system complications.
- Often asymptomatic in early stages, leading to underdiagnosis.
- Disproportionately affects rural, impoverished communities.

Neglect Factors:

- Lack of awareness among affected populations.
- Limited access to diagnostic tools and treatment.
- Insufficient investment in research for better therapies.

Leishmaniasis

Overview: Caused by Leishmania parasites transmitted through sandflies, leishmaniasis manifests in cutaneous, mucocutaneous, and visceral forms, the latter being potentially fatal.

Impact on People:

- Disfiguring skin ulcers affecting self-esteem and social integration.
- Chronic illness leading to malnutrition and economic hardship.
- Predominant in Middle Eastern, African, and South Asian regions.

Neglect Factors:

- Underfunded research.
- Lack of effective, affordable treatments.
- Stigma preventing community engagement.

Schistosomiasis

Overview: Also known as 'snail fever,' caused by parasitic worms (Schistosoma spp.), transmitted through freshwater contaminated with larval forms.

Impact on People:

- Chronic anemia and organ damage.
- Impaired cognitive development in children.
- Reduced productivity among affected populations.

Neglect Factors:

- Often considered a 'disease of poverty' with limited political attention.
- Inadequate sanitation infrastructure perpetuates transmission.

Lymphatic Filariasis (Elephantiasis)

Overview: Caused by filarial worms transmitted via mosquito bites, leading to severe swelling, disfigurement, and disability.

Impact on People:

- Social stigma and exclusion.
- Physical disability impacting livelihoods.
- Psychological distress.

Neglect Factors:

- Lack of awareness.
- Inadequate mass drug administration programs in some regions.

Onchocerciasis (River Blindness)

Overview: Caused by *Onchocerca volvulus* worms transmitted by blackflies, leading to skin disease and blindness.

Impact on People:

- Visual impairment and blindness in affected communities.
- Skin lesions and severe itching.
- Economic consequences due to reduced productivity.

Neglect Factors:

- Historically underfunded, though programs like the African Program for Onchocerciasis Control (APOC) have made progress.

Soil-Transmitted Helminth Infections

Overview: Caused by intestinal worms such as *Ascaris*, *Trichuris*, and hookworms, transmitted through contaminated soil.

Impact on People:

- Malnutrition, anemia, and impaired growth in children.
- Reduced cognitive development.
- Economic burden due to health costs and lost productivity.

Neglect Factors:

- Often overshadowed by more acute diseases.
- Lack of sustained deworming programs in some areas.

Trachoma

Overview: Bacterial infection caused by *Chlamydia trachomatis*, leading to blindness through repeated infections and eyelid scarring.

Impact on People:

- Visual impairment and blindness.
- Social stigma and economic hardship.
- Mainly affects poor communities with limited access to clean water.

Neglect Factors:

- Insufficient funding for mass distribution of antibiotics and sanitation improvements.

Challenges in Addressing Neglected Diseases

Limited Funding and Resources

- Neglected diseases typically receive less than 1% of global health research funding.
- Pharmaceutical companies often lack incentives to develop drugs for diseases prevalent in impoverished regions.

Weak Healthcare Infrastructure

- Many affected regions lack basic healthcare facilities.
- Insufficient diagnostic and treatment capacity hampers disease management.

Social and Cultural Barriers

- Stigma associated with certain diseases discourages treatment-seeking.
- Cultural beliefs may hinder community engagement with health programs.

Political and Policy Neglect

- Limited political will due to competing priorities.
- Neglected diseases rarely feature prominently on national health agendas.

Environmental and Ecological Factors

- Climate change and environmental degradation expand habitats for vectors.
- Urbanization and deforestation influence disease transmission patterns.

Innovative Approaches and Solutions

Integrated Disease Control Programs

- Combining efforts for multiple NTDs in mass drug administration (MDA).
- Improving water, sanitation, and hygiene (WASH) infrastructure.

Advancements in Research and Drug Development

- Development of new, affordable, and effective drugs.
- Use of vector control strategies such as insecticide-treated nets and environmental management.

Community Engagement and Education

- Empowering local populations to participate in disease prevention.
- Culturally sensitive health education campaigns.

Global Partnerships and Funding Initiatives

- WHO's roadmap for NTD elimination.
- Donations from pharmaceutical companies (e.g., Merck's ivermectin donation).
- Funding from global health organizations like the Bill & Melinda Gates Foundation.

Harnessing Technology

- Mobile health (mHealth) for disease surveillance.
- Geographic Information Systems (GIS) for mapping endemic areas.

The Path Forward: Why It Matters

Addressing neglected diseases is not merely a matter of health but also social justice. The communities affected are often among the poorest and most vulnerable, bearing the brunt of both disease and neglect. Their suffering perpetuates cycles of poverty, impeding economic development and social stability.

Key reasons to prioritize neglected diseases:

- Health Equity: Ensuring marginalized populations receive adequate healthcare.
- Global Health Security: Controlling diseases that can spread beyond borders.
- Economic Benefits: Reducing healthcare costs

Question What are 'forgotten diseases' and why are they often overlooked? Forgotten diseases are illnesses that primarily affect impoverished or marginalized populations, receiving little attention or funding for research and treatment. They are often overlooked due to lack of awareness, limited media coverage, and minimal commercial interest. Which diseases are commonly classified as neglected or forgotten diseases? Examples include Chagas disease, leishmaniasis, sleeping sickness, leprosy, and filariasis. These diseases predominantly impact low-income regions and often remain underdiagnosed and undertreated. How do neglected diseases impact local communities and economies? They cause chronic illness, disability, and death, which reduce workforce productivity, increase healthcare costs, and perpetuate cycles of poverty within affected communities. What are some challenges in diagnosing and treating neglected diseases? Challenges include limited healthcare infrastructure, lack of awareness among healthcare providers, insufficient diagnostic tools, and scarce funding for research and development of effective treatments. Why is increased awareness important for neglected diseases? Raising awareness can lead to better funding, research, early diagnosis, and treatment, ultimately reducing

disease burden and improving affected individuals' quality of life. What role do global health organizations play in addressing forgotten diseases? Organizations like WHO and NGOs work to raise awareness, fund research, develop treatment programs, and support healthcare infrastructure in affected regions to combat neglected diseases. How can communities contribute to combating forgotten diseases? Communities can promote health education, support local screening and treatment efforts, advocate for increased funding, and collaborate with health authorities to improve disease awareness and prevention. What recent advancements have been made in the fight against neglected diseases? Recent progress includes the development of new diagnostics, vaccines, and treatments; increased global funding; and successful disease eradication campaigns in certain regions. What is the importance of integrating neglected disease control into broader public health initiatives? Integrating neglected disease efforts into general health programs enhances resource efficiency, improves overall health outcomes, and helps achieve sustainable development goals related to health equity.

Related keywords: marginalized groups, neglected diseases, health disparities, underserved populations, rare diseases, social exclusion, neglected tropical diseases, health inequity, vulnerable communities, disease neglect

Read the full article online: [FORGOTTEN PEOPLE FORGOTTEN DISEASES THE NEGLECTED](#)